

## REVIEW

## Ketone bodies in epilepsy

Melanie A. McNally\* and Adam L. Hartman†

\*Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

†Department of Neurology, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

**Abstract**

Seizures that are resistant to standard medications remain a major clinical problem. One underutilized option for patients with medication-resistant seizures is the high-fat, low-carbohydrate ketogenic diet. The diet received its name based on the observation that patients consuming this diet produce ketone bodies (e.g., acetoacetate,  $\beta$ -hydroxybutyrate, and acetone). Although the exact mechanisms of the diet are unknown, ketone bodies have been hypothesized to contribute to the anticonvulsant and antiepileptic effects. In this review,

anticonvulsant properties of ketone bodies and the ketogenic diet are discussed (including GABAergic and glutamatergic effects). Because of the importance of ketone body metabolism in the early stages of life, the effects of ketone bodies on developing neurons *in vitro* also are discussed. Understanding how ketone bodies exert their effects will help optimize their use in treating epilepsy and other neurological disorders.

**Keywords:**  $\beta$ -hydroxybutyrate, acetoacetate, acetone, epilepsy, ketogenic diet, seizure.

*J. Neurochem.* (2012) 10.1111/j.1471-4159.2012.07670.x

Epilepsy is a devastating and widespread disease that affects around 1% of the US population and at least 50 million people worldwide (Hauser and Kurland 1975; Hirtz *et al.* 2007; Kobau *et al.* 2008). While an array of antiepileptic drugs has been developed, they remain unable to control seizures in approximately 30% of patients with epilepsy (Kwan and Brodie 2000). Some of these patients are candidates for surgical resection of abnormal tissue (e.g., mesial temporal sclerosis) but options are limited for those without such lesions. To help these patients with nonsurgical intractable epilepsy, a high-fat, low-carbohydrate diet known as a ketogenic diet was developed in the 1920s to mimic the beneficial effects of fasting on seizure control (Hartman and Vining 2007). Many clinical studies have confirmed its beneficial effects, with a randomized study showing a 75% decrease in seizures in children on a ketogenic diet for 3 months (Neal *et al.* 2008). However, while many possibilities have been investigated, the mechanisms through which a ketogenic diet confers its anticonvulsant effects remain unknown. Unraveling these mechanisms will allow refinement of its clinical use as well as identification of new targets for future therapy development.


A ketogenic diet receives its name due to the high ketone levels found in the blood and urine of patients consuming the diet, which traditionally consists of either long- or medium-chain fatty acids (Hartman and Vining 2007). In the setting of elevated fatty acids and low dietary carbohydrate content while on a ketogenic diet, the liver produces ketone bodies

by shunting excess acetyl-CoA to ketogenesis (Fig. 1). The two primary ketone bodies produced are acetoacetate and  $\beta$ -hydroxybutyrate. Acetone, the other major ketone elevated in patients on a ketogenic diet, is a metabolite of acetoacetate. During infancy, the brain is much more efficient at extracting and utilizing ketone bodies from the blood due to the higher levels of ketone metabolizing enzymes and monocarboxylic acid transporters produced during this period (Sokoloff 1973; Bilger and Nehlig 1992). Notably, however, the adult brain increases its levels of monocarboxylic acid transporters and ketone metabolizing enzymes rapidly during periods of stress such as ischemia, trauma, or low glucose (Prins 2008). In addition, there is evidence that astrocytes can produce ketone bodies under conditions of glucose deprivation (Blazquez *et al.* 1999). Thus, given the brain's ability to adapt to ketone utilization, the elevated levels of ketone bodies in patients on a ketogenic diet, and the robust anticonvulsant effects observed in these patients, ketone bodies are an attractive candidate to investigate in studies of a ketogenic diet's anticonvulsant mechanism.

Received August 23, 2011; revised manuscript received January 16, 2012; accepted January 16, 2012.

Address correspondence and reprints request to Adam L. Hartman, 600°N. Wolfe St., Meyer 2-147, Baltimore, MD 21287, USA. E-mail: ahartma2@jhmi.edu

*Abbreviations used:* ATP, adenosine tri-phosphate; GABA,  $\gamma$ -aminobutyric acid; VGLUTs, vesicular glutamate transporters.



**Fig. 1** A high-fat, low-carbohydrate ketogenic diet causes a shift in the metabolic activity of hepatocytes. Under these conditions, the Krebs cycle cannot utilize the high levels of acetyl-CoA generated from fat. Remaining acetyl-CoA is converted to the ketone body acetoacetate. The two additional ketone bodies, acetone and  $\beta$ -hydroxybutyrate, are derived from acetoacetate by spontaneous

degradation and enzymatic conversion with  $\beta$ -hydroxybutyrate dehydrogenase, respectively. The three ketone bodies are then released from the hepatocyte, cross the blood-brain barrier, and may exert their effects in the brain. CAT, carnitine-acylcarnitine translocase; ACA, acetoacetate; BHB,  $\beta$ -hydroxybutyrate; BBB, blood-brain barrier.

### Ketone bodies: anticonvulsant properties

In trying to understand a ketogenic diet's therapeutic mechanism, many laboratories have investigated whether ketone bodies at pharmacological doses have intrinsic anticonvulsant properties. The anticonvulsant properties of acetoacetate were first reported by Keith (1935) where its administration in rabbits protected against thujone-induced seizures. In agreement with this early report, Rho and colleagues found that intraperitoneally administered acetoacetate and acetone, but not  $\beta$ -hydroxybutyrate, protected against seizures in the Frings audiogenic seizure-susceptible juvenile mouse (Rho *et al.* 2002). More recently, acute acetone administration by intraperitoneal injection in rats decreased their seizure susceptibility in a broad range of seizure tests including the maximal electroshock test, the subcutaneous pentylenetetrazol test, the amygdala kindling test, and the AY-9944 test (Likhodii *et al.* 2003). Acutely administered acetone, but not its metabolites, protected juvenile mice from seizures induced by pentylenetetrazol and 4-aminopyridine (Gasior *et al.* 2007). However, in this study, the doses of acetone necessary to cause significant protection from seizures caused motor impairment in the mice, limiting its clinical pharmacological implementation. Collectively, the evidence is strong that acutely administered acetone and acetoacetate have intrinsic anticonvulsant properties in many standardized animal models of epilepsy. However, clinical translatability remains unclear. We are unaware of reports examining the anticonvulsant effects of chronic exogenously administered ketone bodies in models similar to those discussed previously.

While acetone has emerged as a broad-spectrum anticonvulsant in animal models, a ketogenic diet in the same tests has not consistently conferred a similar level of protection

(specifically, against pentylenetetrazol-induced seizures in mice) suggesting that acetone and the ketogenic diet may have different anticonvulsant mechanisms (Uhlemann and Neims 1972; Samala *et al.* 2008). Additionally, plasma levels of acetone in rats fed a ketogenic diet were 10-fold lower than the levels found to be therapeutic ( $> 2$  mM), even in the presence of an inhibitor of acetone metabolism (Nylen *et al.* 2006). In contrast, using proton magnetic resonance spectroscopy in epilepsy patients fully controlled by a ketogenic diet, brain acetone levels were approximately 0.7 mM (Seymour *et al.* 1999). These data suggest that humans and rats may produce acetone at different levels in response to a ketogenic diet, possibly explaining the different anticonvulsant profiles observed in animal seizure models using acetone versus a ketogenic diet (Rho *et al.* 2002). To better understand the observed inconsistency (and the possibility that acetone's volatility may lead to inaccurate estimates of actual brain exposure), studies comparing levels of acetone in the brain during a ketogenic diet versus those found with exogenously administered acetone are needed. At this time, acetone's role in the anticonvulsant mechanisms of the ketogenic diet remains unclarified.

### Ketone bodies: possible mechanisms of action

The *in vivo* studies discussed previously suggest that ketone bodies have anticonvulsant properties. A number of theories have emerged to explain the decreased seizure threshold in these models. Glutamate and  $\gamma$ -aminobutyric acid (GABA), respectively, are the major excitatory and inhibitory neurotransmitters in the brain. How ketone bodies affect levels of these neurotransmitters and their receptors, their receptor activity, and their handling has been debated. New theories also are emerging surrounding the effects of ketone bodies on

neuronal membrane potential, neuronal excitability, and reactive oxygen species. Presented here is an overview of the current data evaluating these evolving hypotheses.

### GABA

GABA signaling has remained a key target of investigation since rodent models of epilepsy induced by GABA antagonists have a strong response to a ketogenic diet (Uhlemann and Neims 1972; Bough *et al.* 2002). The role of GABA signaling in the anticonvulsant mechanisms of ketone bodies and/or the ketogenic diet is highly debated. Supporting a role for GABA signaling, GABA levels in rat synaptosomes exposed to acetoacetate or  $\beta$ -hydroxybutyrate are increased (Erecinska *et al.* 1996). More recently, clinical studies of patients on a ketogenic diet show increased GABA levels in the cerebrospinal fluid, as well as with magnetic resonance spectroscopy (Wang *et al.* 2003; Dahlin *et al.* 2005). GABA is synthesized from the Krebs cycle intermediate  $\alpha$ -ketoglutarate, which, via glutamate, is converted to GABA by glutamate decarboxylase. Aspartate is a known inhibitor of glutamate decarboxylase. Therefore, a decrease in aspartate could theoretically promote the synthesis of GABA. Exploring this further, a 1-h exposure to acetoacetate or  $\beta$ -hydroxybutyrate was found to decrease aspartate levels in cultured astrocytes (Yudkoff *et al.* 1997). A similar decrease in aspartate levels was detected in the forebrain and cerebellum of mice fed a ketogenic diet (Yudkoff *et al.* 2001). The body of data from Yudkoff's group suggest a model wherein the ketotic brain has increased conversion of glutamate to glutamine in astrocytes; glutamine is then taken up into neurons, where it is then converted eventually to GABA (preferentially over aspartate), thus increasing neuronal inhibition (Yudkoff *et al.* 2008). One unanswered question about the importance of GABA is why a ketogenic diet did not protect murine models of epilepsy induced by the GABA antagonist pentylenetetrazol in the same way it did rats (Uhlemann and Neims 1972). Furthermore, a number of studies have shown no changes in GABA levels from whole brain samples of rodents fed a ketogenic diet, seemingly inconsistent with an increase in GABA synthesis (Appleton and DeVivo 1974; Al-Mudallal *et al.* 1996; Yudkoff *et al.* 2001, 2005; Melo *et al.* 2006). Additional studies are required to investigate whether there are local or regional changes in GABA levels, GABA shunt flow, or GABA responsiveness within the brain in the presence of ketone bodies.

### VGLUTs

Ketone bodies may alter the behavior of vesicular glutamate transporters (VGLUTs) that are responsible for filling pre-synaptic vesicles with glutamate in a  $\text{Cl}^-$ -dependent manner. Using proteoliposomes as an *in vitro* model, Juge and colleagues demonstrated that  $\text{Cl}^-$  is an allosteric activator of VGLUTs that is competitively inhibited by ketone bodies, acetoacetate more strongly than  $\beta$ -hydroxybutyrate (Juge

*et al.* 2010). Strengthening their *in vitro* findings, they showed that acetoacetate reversibly inhibited glutamate release from both cultured rat neurons and mouse CA1 pyramidal neurons in hippocampal slices. Finally, acetoacetate was able to suppress *in vivo* glutamate release and seizures in rat brains exposed to 4-aminopyridine. However, the amount of acetoacetate that the authors used (10 mM) is likely supratherapeutic, given that hippocampal extracellular levels of  $\beta$ -hydroxybutyrate are 40–50  $\mu\text{M}$  in mice fed a ketogenic diet for 2–3 weeks (Samala *et al.* 2011). As these two compounds have a stoichiometric relationship (Owen *et al.* 1967), it is expected that levels of acetoacetate would be significantly < 10 mM in these mice. In addition, children on a ketogenic diet have been found to have cerebrospinal fluid levels of acetoacetate at 0.3 mM, much lower than the concentrations utilized in the Juge *et al.* study (Nordli and de Vivo 1997). Therefore, the clinical relevance of higher levels used by Juge *et al.* needs to be further clarified.

### Other neurotransmitters, transporters, and receptors

Controversy remains over how ketone bodies affect neurotransmitter responsiveness. Available data suggest that a ketogenic diet does not alter levels of the neuronal glutamate transporter EAAC1 in neocortex, hippocampus, or cerebellum, measured by Western blot, electron microscopy and glutamate uptake in hippocampal proteoliposomes (Bough *et al.* 2007). *In vitro* patch clamp studies of post-synaptic GABA<sub>A</sub> receptors on rat hippocampal neurons suggest that there is no change in receptor activity in the presence of extracellular 2 mM  $\beta$ -hydroxybutyrate or 1 mM acetoacetate (Thio *et al.* 2000). In these studies, the ketone body containing solutions, which also contained glucose, were pre-incubated with the neurons for only 1 min. Examining the behavior of the receptors following longer-term incubation with ketone bodies in the absence of glucose would be of interest as these conditions model more closely the clinical scenario in patients on a ketogenic diet. More recent work using *Xenopus* oocytes expressing human GABA<sub>A</sub> receptors shows that a 5-min incubation with 10 mM  $\beta$ -hydroxybutyrate or 50 mM acetone significantly increases GABA<sub>A</sub> receptor activity (Yang *et al.* 2007). While intriguing, Yang and colleagues did not find similar changes at lower concentrations of the ketone bodies that would be seen clinically. Good *in vivo* data to clarify what role, if any, that altered GABA receptor responsiveness plays in the anticonvulsant activity of ketone bodies, are lacking.

Other neurotransmitter systems not typically considered in epilepsy also appear to be necessary for the ketogenic diet's anticonvulsant effects. Data for an effect of norepinephrine includes the lack of a ketogenic diet's protective effects against the convulsant fluorothyl in dopamine  $\beta$ -hydroxylase knockout mice that cannot synthesize norepinephrine (Sztot *et al.* 2001). Evidence of the need for adenosine neurotransmission was shown when mice with adenosine A<sub>1</sub> receptor mutations were not protected against recurrent seizures after

ketogenic diet treatment (Masino *et al.* 2011). A ketogenic diet also decreased the adenosine metabolizing enzyme, adenosine kinase. Other neurotransmitters have not been investigated as thoroughly.

### *K<sub>ATP</sub> channels*

Another way ketone bodies may be anticonvulsant is through a direct effect on neuronal membrane potential. K<sub>ATP</sub> channels are of particular interest given their close relationship with cellular metabolism [activated when cellular adenosine tri-phosphate (ATP) is low] and their hyperpolarizing effects. Additionally, they were found to play a key role in CA3 hippocampal neuron hyperpolarization when the neurons were cultured in low glucose media (Kawamura *et al.* 2010). Exploring this idea further, in cultured mouse hippocampal dentate gyrus neurons, both basal levels of inward-rectifying K<sub>ATP</sub> channel opening and the amount of stimulus evoked K<sub>ATP</sub> channel opening was increased in the presence of β-hydroxybutyrate (Tanner *et al.* 2011). Adding support to this *in vitro* work, live recordings of the hippocampal dentate gyrus in rats fed a ketogenic diet show decreased long-term potentiation, consistent with increased neuronal inhibition (Koranda *et al.* 2011). Previously, Ma and colleagues used GABAergic substantia nigra pars reticulata neurons to study K<sub>ATP</sub> channels given the hypothesis that very rapid firing of these neurons precede generalized seizure onset (Deransart *et al.* 2003; Ma *et al.* 2007). When GABAergic substantia nigra pars reticulata neurons were cultured in the presence of acetoacetate or β-hydroxybutyrate with glucose, firing rates (measured by loose-patch cell-attached recordings) were decreased. Of note, this effect was abolished in a variety of settings, including (a) when the neurons were cultured from Kir6.2 knockout mice (no cell surface K<sub>ATP</sub> channels), and (b) when the experiments were conducted in the presence of tolbutamide (an inhibitor of K<sub>ATP</sub> channel opening) or a GABA<sub>B</sub> receptor blocker. These findings add further support to the hypothesis that the anticonvulsant activities of ketone bodies are, in part, mediated by their effects on K<sub>ATP</sub> channels. The latter scenario suggests that GABA<sub>B</sub> signaling might be important to mediate these effects of ketone bodies. However, whether these results can shed light on mechanisms of the ketogenic diet needs to be considered carefully as rodent studies to date have shown that a ketogenic diet elevates or maintains normal cellular ATP levels, an otherwise negative regulator of K<sub>ATP</sub> channels (DeVivo *et al.* 1978; Bough *et al.* 2006; Nylen *et al.* 2009). The notion has been advanced that a ketogenic diet enhances K<sub>ATP</sub> channel activity through ATP compartmentation by decreasing glycolytic ATP levels, an hypothesis that awaits further experimental evidence (Yellen 2008).

### *Krebs cycle and the electron transport chain*

One theory of the ketogenic diet's multifaceted effects is that it may optimize Krebs cycle function, with a subsequent

effect on neuronal function. The logical starting point for this theory is that acetyl CoA (one downstream product of ketogenesis) is a primary substrate for the Krebs cycle (Fig. 1). Downstream of the Krebs cycle, β-hydroxybutyrate improved redox activity and decreased cell death in dopaminergic SH-SY5Y cells after exposure to the complex I toxin rotenone (Imamura *et al.* 2006). β-Hydroxybutyrate appears to require an intact complex II to confer its beneficial effects in two different models of neurodegeneration, including a complex II toxin (3-nitropropionic acid) model and the SOD1-G93A mouse model of amyotrophic lateral sclerosis (Lou Gehrig's disease) (Tieu *et al.* 2003; Zhao *et al.* 2006). More recently, Kim do *et al.* (2010) showed in hippocampal slices that a cocktail of β-hydroxybutyrate and acetoacetate preserved synaptic activity after exposure to rotenone or 3-nitropropionic acid, although there were some regional disparities (i.e., different responses even within the same slice) noted during ketone exposure to the latter. The reason for differences between the two studies of 3-nitropropionic acid is unclear.

A ketogenic diet also has been shown to induce transcription of some electron transport chain subunit mRNAs (Bough *et al.* 2006). An increased number of electron transport chain subunits might increase neuronal 'energy reserves' and, in fact, ketogenic diets increase brain ATP levels in some (but not all) rodent studies (DeVivo *et al.* 1978; Bough *et al.* 2006; Nylen *et al.* 2009; Kim do *et al.* 2010). A final hypothesis is that increased ATP levels may stabilize neuronal membrane potential by enhancing the function of Na<sup>+</sup>-K<sup>+</sup>-ATPases (Bough and Rho 2007). Although teleologically satisfying, the events outlined here have not been completely linked experimentally in either normal or epilepsy animal models. Nonetheless, individual experiments point to the potential importance of the Krebs cycle and electron transport chain function in the effects of ketone bodies.

### *Immature and developing neurons*

Many studies have looked at the effects of ketone bodies on immature neurons as a possible physiological model of the ketogenic diet. In suckling rodents, circulating levels of ketone bodies are high due to the high ketogenic ratio of rodent milk (Nehlig 1999). During this time, the brain relies most heavily on ketone bodies as a source of metabolic fuel (Nehlig and Pereira de Vasconcelos 1993). Also during this time, GABA is excitatory, an action found to be dependent on the higher intracellular chloride levels in immature neurons. As a neuron matures the Na<sup>+</sup>-K<sup>+</sup>-2Cl<sup>-</sup> co-transporter (NKCC1) is replaced with the K<sup>+</sup>-Cl<sup>-</sup> co-transporter (KCC2) shifting GABA to an inhibitory neurotransmitter (Aronica *et al.* 2011). Patch clamp work on neocortical pyramidal neurons from mice demonstrated that β-hydroxybutyrate incubation at supra-physiologic levels decreased both the resting membrane potential and the amount of GABA-mediated excitation in a dose-dependent manner

(Rheims *et al.* 2009). This group also showed that ketone bodies at similarly high levels cause the same inhibition of GABA-mediated excitation in murine hippocampal CA3 neurons (Holmgren *et al.* 2010), another important brain region in epilepsy development. These data are supported by recent *in vivo* work examining physiological levels of ketone bodies in neonatal rats (Minlebaev and Khazipov 2011). Minlebaev and Khazipov used 16-channel extracellular field potential recordings from somatosensory barrel cortex in postnatal day 5–9 awake rat pups to study epileptiform activity induced by flurothyl when physiological ketosis was inhibited. They found that blocking ketogenesis did not affect a single flurothyl-evoked seizure, however consecutively evoked seizures were more severe in these animals. In contrast to the preceding studies, a number of groups have found that GABA-induced depolarization in neonatal rodent cortical neurons is unaffected by the presence of ketone bodies at both supraphysiologic and physiologic levels (Thio *et al.* 2000; Kirmse *et al.* 2010; Tyzio *et al.* 2011). Possible reasons for the discrepancies among the studies are outlined in Tyzio *et al.* (2011). Thus, while an effect of ketone bodies on neonatal cortex excitability remains an intriguing possibility that could shed light on possible therapeutic mechanisms of a ketogenic diet, its existence remains controversial. These data also should be interpreted with caution when drawing parallels to a ketogenic diet as all these results were found with solutions containing both ketone bodies and high levels of glucose (5–10 mM). Basal glucose concentrations

in murine hippocampal extracellular fluid as determined by *in vivo* microdialysis were approximately 2 mM, and only ~1.3 mM when the mice were fed a ketogenic diet (Samala *et al.* 2011).


#### Antioxidant effects

Ketone bodies also may have antioxidant effects. Reactive oxygen species production resulting from glutamate exposure is inhibited by ketone bodies in primary cultures of rat neocortical neurons (Maalouf *et al.* 2007). The mechanism appears to involve NADH oxidation, rather than a glutathione-mediated antioxidant effect. Conversely, the ketogenic diet (rather than ketone bodies) may alter glutathione metabolism via the transcription factor Nrf2 (Milder *et al.* 2010). Ketone bodies also protect against cell death after neocortical slices are exposed to hydrogen peroxide (Kim do *et al.* 2007). Additionally, ketone bodies prevent hydrogen peroxide-induced impairment of hippocampal long-term potentiation (Maalouf and Rho 2008). Therefore, in addition to effects on neurotransmission, these data suggest ketone bodies may be neuroprotective against reactive oxygen species produced in neuropathological states.

#### Conclusions

A growing amount of *in vivo* and *in vitro* data show that ketone bodies can alter neuronal activity and protect against

**Fig. 2** Possible anticonvulsant effects of ketone bodies on the brain. (1) Increased GABA synthesis through alteration of glutamate cycling in glutamate-glutamine cycle or altered neuronal responsiveness to GABA at GABA<sub>A</sub> receptors. (2) Decreased glutamate release by competitive inhibition of vesicular glutamate transporters. (3) Other neurotransmitters, including norepinephrine and adenosine. (4) Increased membrane potential hyperpolarization via K<sub>ATP</sub> channels possibly mediated by GABA<sub>B</sub> receptor signaling. (5) Decreased reactive oxygen species production from glutamate exposure. (6) Electron transport chain subunit transcription. A<sub>1</sub>R, adenosine receptor; Cl<sup>-</sup>, chloride; GLN, glutamine; GLU, glutamate; GABA,  $\gamma$ -aminobutyric acid; GABA<sub>B</sub>R,  $\gamma$ -aminobutyric acid beta receptor; GABA<sub>A</sub>R,  $\gamma$ -aminobutyric acid alpha receptor; VGLUT, vesicular glutamate transporter; ROS, reactive oxygen species.



seizures. The mechanisms through which ketone bodies may confer an anticonvulsant effect are still debated. The potential mechanisms discussed in this review center around neurotransmitters and the resting neuronal membrane potential (Fig. 2). Moving forward, work will need to be completed exploring how physiological levels of ketone bodies affect these systems. The exact targets of ketone bodies still are unknown and may be direct (i.e., binding to a specific neuronal target, as demonstrated for acetoacetate in VGLUT), or indirect (i.e., supplying carbon equivalents to the Krebs cycle or an antioxidant effect). Given both the success of AC-1202 (an oral ketogenic compound that rapidly elevates  $\beta$ -hydroxybutyrate levels) in improving cognitive outcomes in Alzheimer disease patients (Henderson *et al.* 2009) and the success of a ketogenic diet in treating children with medically intractable epilepsy, there is significant interest in pursuing these types of compounds for epilepsy therapies. However, the contribution of ketone bodies to the anticonvulsant effects of the ketogenic diet remains elusive. Because of study design issues and the amount of overlap between values in different patient groups, the relationship between seizure control and serum levels of  $\beta$ -hydroxybutyrate remains unclear (Gilbert *et al.* 2000). Many other studies have failed to show that the levels of serum or urine ketosis correlate with seizure protection in animal models (Bough *et al.* 2000; Likhodii *et al.* 2000) or patients (Ross *et al.* 1985; Fraser *et al.* 2003). Breath acetone levels correlate with plasma levels of all three ketone bodies but not with seizure control (Musa-Veloso *et al.* 2006). Some of these studies highlight the potential difference between administration of ketone bodies versus a ketogenic diet (understanding the former as a product of the latter). Together, some of the data discussed here suggest the ketogenic diet may have anticonvulsant effects via mechanisms other than elevated ketone body levels, even though ketone bodies have numerous potential beneficial effects. Therefore, based on current data, ketone bodies may represent a unique, additional therapeutic modality to optimize for patients with epilepsy and possibly other neurological disorders. Further studies are required to understand how using these compounds may help these populations.

## Acknowledgements

Supported by the Howard Hughes Medical Institute Medical Research Fellows Program (MAM), K12NS001696, K08NS070931, and a Clinician Scientist Award (Johns Hopkins University School of Medicine) (ALH). The authors have no conflicts of interest to disclose.

## References

- Al-Mudallal A. S., LaManna J. C., Lust W. D. and Harik S. I. (1996) Diet-induced ketosis does not cause cerebral acidosis. *Epilepsia* **37**, 258–261.
- Appleton D. B. and DeVivo D. C. (1974) An animal model for the ketogenic diet. *Epilepsia* **15**, 211–227.
- Aronica E., Iyer A., Zurolo E. and Gorter J. A. (2011) Ontogenetic modifications of neuronal excitability during brain maturation: developmental changes of neurotransmitter receptors. *Epilepsia* **52** (Suppl. 8), 3–5.
- Bilger A. and Nehlig A. (1992) Quantitative histochemical changes in enzymes involved in energy metabolism in the rat brain during postnatal development. II. Glucose-6-phosphate dehydrogenase and beta-hydroxybutyrate dehydrogenase. *Int. J. Dev. Neurosci.* **10**, 143–152.
- Blazquez C., Woods A., de Ceballos M. L., Carling D. and Guzman M. (1999) The AMP-activated protein kinase is involved in the regulation of ketone body production by astrocytes. *J. Neurochem.* **73**, 1674–1682.
- Bough K. J. and Rho J. M. (2007) Anticonvulsant mechanisms of the ketogenic diet. *Epilepsia* **48**, 43–58.
- Bough K. J., Yao S. G. and Eagles D. A. (2000) Higher ketogenic diet ratios confer protection from seizures without neurotoxicity. *Epilepsy Res.* **38**, 15–25.
- Bough K. J., Gudi K., Han F. T., Rathod A. H. and Eagles D. A. (2002) An anticonvulsant profile of the ketogenic diet in the rat. *Epilepsy Res.* **50**, 313–325.
- Bough K. J., Wetherington J., Hassel B. *et al.* (2006) Mitochondrial biogenesis in the anticonvulsant mechanism of the ketogenic diet. *Ann. Neurol.* **60**, 223–235.
- Bough K. J., Paquet M., Pare J. F., Hassel B., Smith Y., Hall R. A. and Dingledine R. (2007) Evidence against enhanced glutamate transport in the anticonvulsant mechanism of the ketogenic diet. *Epilepsy Res.* **74**, 232–236.
- Dahlin M., Elfving A., Ungerstedt U. and Amark P. (2005) The ketogenic diet influences the levels of excitatory and inhibitory amino acids in the CSF in children with refractory epilepsy. *Epilepsy Res.* **64**, 115–125.
- Deransart C., Hellwig B., Heupel-Reuter M., Leger J. F., Heck D. and Lucking C. H. (2003) Single-unit analysis of substantia nigra pars reticulata neurons in freely behaving rats with genetic absence epilepsy. *Epilepsia* **44**, 1513–1520.
- DeVivo D. C., Leckie M. P., Ferrendelli J. S. and McDougal D. B. Jr (1978) Chronic ketosis and cerebral metabolism. *Ann. Neurol.* **3**, 331–337.
- Erecinska M., Nelson D., Daikhin Y. and Yudkoff M. (1996) Regulation of GABA level in rat brain synaptosomes: fluxes through enzymes of the GABA shunt and effects of glutamate, calcium, and ketone bodies. *J. Neurochem.* **67**, 2325–2334.
- Fraser D. D., Whiting S., Andrew R. D., Macdonald E. A., Musa-Veloso K. and Cunnane S. C. (2003) Elevated polyunsaturated fatty acids in blood serum obtained from children on the ketogenic diet. *Neurology* **60**, 1026–1029.
- Gasior M., French A., Joy M. T., Tang R. S., Hartman A. L. and Rogawski M. A. (2007) The anticonvulsant activity of acetone, the major ketone body in the ketogenic diet, is not dependent on its metabolites acetol, 1,2-propanediol, methylglyoxal, or pyruvic acid. *Epilepsia* **48**, 793–800.
- Gilbert D. L., Pyzik P. L. and Freeman J. M. (2000) The ketogenic diet: seizure control correlates better with serum beta-hydroxybutyrate than with urine ketones. *J. Child Neurol.* **15**, 787–790.
- Hartman A. L. and Vining E. P. (2007) Clinical aspects of the ketogenic diet. *Epilepsia* **48**, 31–42.
- Hauser W. A. and Kurland L. T. (1975) The epidemiology of epilepsy in Rochester, Minnesota, 1935 through 1967. *Epilepsia* **16**, 1–66.
- Henderson S. T., Vogel J. L., Barr L. J., Garvin F., Jones J. J. and Costantini L. C. (2009) Study of the ketogenic agent AC-1202 in mild to moderate Alzheimer's disease: a randomized, double-

- blind, placebo-controlled, multicenter trial. *Nutr. Metab. (Lond)* **6**, 31.
- Hirtz D., Thurman D. J., Gwinn-Hardy K., Mohamed M., Chaudhuri A. R. and Zalutsky R. (2007) How common are the “common” neurologic disorders? *Neurology* **68**, 326–337.
- Holmgren C. D., Mukhtarov M., Malkov A. E., Popova I. Y., Bregestovski P. and Zilberter Y. (2010) Energy substrate availability as a determinant of neuronal resting potential, GABA signaling and spontaneous network activity in the neonatal cortex *in vitro*. *J. Neurochem.* **112**, 900–912.
- Imamura K., Takeshima T., Kashiwaya Y., Nakaso K. and Nakashima K. (2006) D-beta-hydroxybutyrate protects dopaminergic SH-SY5Y cells in a rotenone model of Parkinson’s disease. *J. Neurosci. Res.* **84**, 1376–1384.
- Juge N., Gray J. A., Omote H. *et al.* (2010) Metabolic control of vesicular glutamate transport and release. *Neuron* **68**, 99–112.
- Kawamura M. Jr, Ruskin D. N. and Masino S. A. (2010) Metabolic autocrine regulation of neurons involves cooperation among pannexin hemichannels, adenosine receptors, and KATP channels. *J. Neurosci.* **30**, 3886–3895.
- Keith H. (1935) Experimental convulsions induced by administration of thujone. *Arch. Neurol. Psychiatry* **34**, 1022–1040.
- Kim do Y., Davis L. M., Sullivan P. G., Maalouf M., Simeone T. A., van Brederode J. and Rho J. M. (2007) Ketone bodies are protective against oxidative stress in neocortical neurons. *J. Neurochem.* **101**, 1316–1326.
- Kim do Y., Vallejo J. and Rho J. M. (2010) Ketones prevent synaptic dysfunction induced by mitochondrial respiratory complex inhibitors. *J. Neurochem.* **114**, 130–141.
- Kirmse K., Witte O. W. and Holthoff K. (2010) GABA depolarizes immature neocortical neurons in the presence of the ketone body ss-hydroxybutyrate. *J. Neurosci.* **30**, 16002–16007.
- Kobay R., Zahran H., Thurman D. J., Zack M. M., Henry T. R., Schachter S. C. and Price P. H. (2008) Epilepsy surveillance among adults—19 states, behavioral risk factor surveillance system, 2005. *MMWR Surveill. Summ.* **57**, 1–20.
- Koranda J. L., Ruskin D. N., Masino S. A. and Blaise J. H. (2011) A ketogenic diet reduces long-term potentiation in the dentate gyrus of freely behaving rats. *J. Neurophysiol.* **106**, 662–666.
- Kwan P. and Brodie M. J. (2000) Epilepsy after the first drug fails: substitution or add-on? *Seizure* **9**, 464–468.
- Likhodii S. S., Musa K., Mendonca A., Dell C., Burnham W. M. and Cunnane S. C. (2000) Dietary fat, ketosis, and seizure resistance in rats on the ketogenic diet. *Epilepsia* **41**, 1400–1410.
- Likhodii S. S., Serbanescu I., Cortez M. A., Murphy P., Snead O. C. 3rd, and Burnham W. M. (2003) Anticonvulsant properties of acetone, a brain ketone elevated by the ketogenic diet. *Ann. Neurol.* **54**, 219–226.
- Ma W., Berg J. and Yellen G. (2007) Ketogenic diet metabolites reduce firing in central neurons by opening K(ATP) channels. *J. Neurosci.* **27**, 3618–3625.
- Maalouf M. and Rho J. M. (2008) Oxidative impairment of hippocampal long-term potentiation involves activation of protein phosphatase 2A and is prevented by ketone bodies. *J. Neurosci. Res.* **86**, 3322–3330.
- Maalouf M., Sullivan P. G., Davis L., Kim D. Y. and Rho J. M. (2007) Ketones inhibit mitochondrial production of reactive oxygen species production following glutamate excitotoxicity by increasing NADH oxidation. *Neuroscience* **145**, 256–264.
- Masino S. A., Li T., Theofilas P., Sandau U. S., Ruskin D. N., Fredholm B. B., Geiger J. D., Aronica E. and Boison D. (2011) A ketogenic diet suppresses seizures in mice through adenosine A receptors. *J. Clin. Invest.* **121**, 2679–2683.
- Melo T. M., Nehlig A. and Sonnewald U. (2006) Neuronal-glia interactions in rats fed a ketogenic diet. *Neurochem. Int.* **48**, 498–507.
- Milder J. B., Liang L. P. and Patel M. (2010) Acute oxidative stress and systemic Nrf2 activation by the ketogenic diet. *Neurobiol. Dis.* **40**, 238–244.
- Minlebaev M. and Khazipov R. (2011) Antiepileptic effects of endogenous beta-hydroxybutyrate in suckling infant rats. *Epilepsy Res.* **95**, 100–109.
- Musa-Veloso K., Likhodii S. S., Rarama E. *et al.* (2006) Breath acetone predicts plasma ketone bodies in children with epilepsy on a ketogenic diet. *Nutrition* **22**, 1–8.
- Neal E. G., Chaffe H., Schwartz R. H., Lawson M. S., Edwards N., Fitzsimmons G., Whitney A. and Cross J. H. (2008) The ketogenic diet for the treatment of childhood epilepsy: a randomised controlled trial. *Lancet Neurol.* **7**, 500–506.
- Nehlig A. (1999) Age-dependent pathways of brain energy metabolism: the suckling rat, a natural model of the ketogenic diet. *Epilepsy Res.* **37**, 211–221.
- Nehlig A. and Pereira de Vasconcelos A. (1993) Glucose and ketone body utilization by the brain of neonatal rats. *Prog. Neurobiol.* **40**, 163–221.
- Nordli D. R. Jr and de Vivo D. C. (1997) The ketogenic diet revisited: back to the future. *Epilepsia* **38**, 743–749.
- Nylen K., Likhodii S. S., Hum K. M. and Burnham W. M. (2006) A ketogenic diet and diallyl sulfide do not elevate afterdischarge thresholds in adult kindled rats. *Epilepsy Res.* **71**, 23–31.
- Nylen K., Velazquez J. L., Sayed V., Gibson K. M., Burnham W. M., and Snead O. C. 3rd (2009) The effects of a ketogenic diet on ATP concentrations and the number of hippocampal mitochondria in Aldh5a1(-/-) mice. *Biochim. Biophys. Acta* **1790**, 208–212.
- Owen O. E., Morgan A. P., Kemp H. G., Sullivan J. M., Herrera M. G. and Cahill G. F. Jr (1967) Brain metabolism during fasting. *J. Clin. Invest.* **46**, 1589–1595.
- Prins M. L. (2008) Cerebral metabolic adaptation and ketone metabolism after brain injury. *J. Cereb. Blood Flow Metab.* **28**, 1–16.
- Rheims S., Holmgren C. D., Chazal G., Mulder J., Harkany T., Zilberter T. and Zilberter Y. (2009) GABA action in immature neocortical neurons directly depends on the availability of ketone bodies. *J. Neurochem.* **110**, 1330–1338.
- Rho J. M., Anderson G. D., Donevan S. D. and White H. S. (2002) Acetoacetate, acetone, and dibenzylamine (a contaminant in l-(+)-beta-hydroxybutyrate) exhibit direct anticonvulsant actions *in vivo*. *Epilepsia* **43**, 358–361.
- Ross D. L., Swaiman K. F., Torres F. and Hansen J. (1985) Early biochemical and EEG correlates of the ketogenic diet in children with atypical absence epilepsy. *Pediatr. Neurol.* **1**, 104–108.
- Samala R., Willis S. and Borges K. (2008) Anticonvulsant profile of a balanced ketogenic diet in acute mouse seizure models. *Epilepsy Res.* **81**, 119–127.
- Samala R., Klein J. and Borges K. (2011) The ketogenic diet changes metabolite levels in hippocampal extracellular fluid. *Neurochem. Int.* **58**, 5–8.
- Seymour K. J., Bluml S., Sutherland J., Sutherland W. and Ross B. D. (1999) Identification of cerebral acetone by 1H-MRS in patients with epilepsy controlled by ketogenic diet. *MAGMA* **8**, 33–42.
- Sokoloff L. (1973) Metabolism of ketone bodies by the brain. *Annu. Rev. Med.* **24**, 271–280.
- Szot P., Weinschenker D., Rho J. M., Storey T. W. and Schwartzkroin P. A. (2001) Norepinephrine is required for the anticonvulsant effect of the ketogenic diet. *Brain Res. Dev. Brain Res.* **129**, 211–214.
- Tanner G. R., Lutas A., Martinez-Francois J. R. and Yellen G. (2011) Single K ATP channel opening in response to action potential firing in mouse dentate granule neurons. *J. Neurosci.* **31**, 8689–8696.

- Thio L. L., Wong M. and Yamada K. A. (2000) Ketone bodies do not directly alter excitatory or inhibitory hippocampal synaptic transmission. *Neurology* **54**, 325–331.
- Tieu K., Perier C., Caspersen C. *et al.* (2003) D-beta-hydroxybutyrate rescues mitochondrial respiration and mitigates features of Parkinson disease. *J. Clin. Invest.* **112**, 892–901.
- Tyzio R., Allene C., Nardou R. *et al.* (2011) Depolarizing actions of GABA in immature neurons depend neither on ketone bodies nor on pyruvate. *J. Neurosci.* **31**, 34–45.
- Uhlemann E. R. and Neims A. H. (1972) Anticonvulsant properties of the ketogenic diet in mice. *JPET* **180**, 231–238.
- Wang Z. J., Bergqvist C., Hunter J. V., Jin D., Wang D. J., Wehrli S. and Zimmerman R. A. (2003) *In vivo* measurement of brain metabolites using two-dimensional double-quantum MR spectroscopy—exploration of GABA levels in a ketogenic diet. *Magn. Reson. Med.* **49**, 615–619.
- Yang L., Zhao J., Milutinovic P. S., Brosnan R. J., Eger E. I. 2nd, and Sonner J. M. (2007) Anesthetic properties of the ketone bodies beta-hydroxybutyric acid and acetone. *Anesth. Analg.* **105**, 673–679.
- Yellen G. (2008) Ketone bodies, glycolysis, and KATP channels in the mechanism of the ketogenic diet. *Epilepsia* **49** (Suppl. 8), 80–82.
- Yudkoff M., Daikhin Y., Nissim I. and Grunstein R. (1997) Effects of ketone bodies on astrocyte amino acid metabolism. *J. Neurochem.* **69**, 682–692.
- Yudkoff M., Daikhin Y., Nissim I. and Lazarow A. (2001) Brain amino acid metabolism and ketosis. *J. Neurosci. Res.* **66**, 272–281.
- Yudkoff M., Daikhin Y., Nissim I., Horyn O., Lazarow A., Luhovyy B. and Wehrli S. (2005) Response of brain amino acid metabolism to ketosis. *Neurochem. Int.* **47**, 119–128.
- Yudkoff M., Daikhin Y., Horyn O. and Nissim I. (2008) Ketosis and brain handling of glutamate, glutamine, and GABA. *Epilepsia* **49** (Suppl. 8), 73–75.
- Zhao Z., Lange D. J., Voustantiouk A. *et al.* (2006) A ketogenic diet as a potential novel therapeutic intervention in amyotrophic lateral sclerosis. *BMC Neurosci.* **7**, 29.